

# Nurses' Knowledge and Attitudes Toward Deep Vein Thrombosis: A Cross-Sectional Study

Zainab Salman Dawood, MSc\* Khadija Mohammed Jassim, MSc\*\* Ali Malik Tiryag, MSc\*\* Abdulkareem Salman Khudhair, PhD\*\*

## ABSTRACT

**Aim:** To ascertain the relationship between the nurses' socio-demographic data (age, gender, level of education, place of employment, and years of experience) and their knowledge about and attitudes toward deep vein thrombosis.

**Methods:** An analysis of the attitudes and information of nurses regarding deep vein thrombosis that is descriptive (cross-sectional). The study was carried out in Al-Basrah's teaching hospitals. The study period's beginning date was changed from August 1st, 2022, to March 1st, 2023. a non-probability (purposive) sample of 100 nurses. A pilot study involved ten nurses who worked at Al-Basrah Teaching Hospital between September 6 and September 21. Data were acquired through developing the questionnaire. When analyzing descriptive data, the mean of the score, the standard deviation (SD), and the frequency are all taken into consideration.

**Findings:** Just 2% of nurses had good awareness of deep vein thrombosis, while 22% of nurses had intermediate knowledge and 76% had inadequate knowledge of the condition. The study's findings also revealed that (83%) of nurses have negative attitudes of deep vein thrombosis, (13%) have moderate understanding, and (4%), have positive attitudes.

**Conclusion:** The current study found that nurses' attitudes and knowledge of deep vein thrombosis were low.

**Keywords:** Nurses, Knowledge, Attitudes, Deep Vein Thrombosis

## INTRODUCTION

Patients frequently get deep vein thrombosis, which can have disastrous effects. In order to reduce problems, nurses' knowledge and knowledge of DVT dangers and prophylaxis must be improved. According to the World Health Organization (WHO), non-communicable disease mortality is increasing globally, with emerging nations accounting for up to 80% of these fatalities. After heart attack and stroke, venous thromboembolism (VTE) is the third most common vascular diagnosis. Obesity and metabolic disorders including diabetes mellitus, all of which increase the risk of having DVT/VTE, are on the rise<sup>1</sup>. Due to its close association with a number of leading causes of death, including as heart disease, type 2 diabetes, and several malignancies<sup>2</sup>, obesity is a global health concern. Extreme obesity can be successfully treated with bariatric surgery, which results in sustained weight loss over time, an improvement in quality of life, and increased longevity<sup>3</sup>. It also reduces and reverses a number of obesity-related comorbidities. One of the most dangerous postoperative complications of TKA is deep vein thrombosis (DVT), which can result in pulmonary embolism (PE) and death<sup>4-7</sup>. In comparison to total hip arthroplasty (THA), TKA is linked to a higher risk of DVT<sup>8,9</sup>. Numerous studies<sup>10,11</sup> have shown the connection between the state of inflammation and hypercoagulability. The past ten years have seen a lot of research on inflammatory markers that are predictive in the diagnosis of post-TKA DVT<sup>12-14</sup>. One of the better researched inflammatory markers for predicting problems with TKA is interleukin-6 (IL-6), which has been compared to C-reactive protein (CRP)<sup>15-19</sup>. Deep vein thrombosis is a common and fatal disorder that can be avoided and managed, and as a result, the mortality and morbidity rates of patients increase<sup>20</sup>. Deep vein thrombosis (DVT) and pulmonary embolism (PE) are both components of VTE, which

has affected more than 900,000 people in the USA. Each year, between 60,000 and 100,000 of these people pass away from VTE<sup>21,22</sup>. Deep vein thrombosis, which is the third most common cardiovascular condition after myocardial infarction and stroke with a recurrence rate of 26.4% of cases, is a serious public health issue. This raises the expense of patient care and the burden on hospitals<sup>23</sup>. Between 3% and 28% more people have DVT in Asia than there were ten years ago. In the past 30 years, the prevalence of VTE has nearly increased by six, especially as an orthopedic surgery complication<sup>24-26</sup>. Distal DVT may return or spread to the proximal veins, especially if untreated, which raises the possibility of complications including PE and PTS. Even though it is uncommon, symptomatic PE may manifest as a right heart overload, tachycardia, discomfort on inspiration, and shortness of breath as a side effect of isolated distal DVT. It may cause circulatory collapse and death if left untreated. People with isolated DVT had a lower yearly rate of VTE recurrence after anticoagulation than those with proximal DVT, but a comparable risk of PE recurrence<sup>27</sup>.

## INSTRUMENT AND METHODS

An examination of nurses' attitudes toward and knowledge of deep vein thrombosis in a cross-sectional descriptive research. "Al-Zubair Hospital, Al-Mawani Teaching Hospital, Al-Basrah Teaching Hospital, Al-Sader Teaching Hospital, and Al-Faiha Teaching Hospital were the Al-Basrah Teaching Hospitals" where the study was conducted. The start of the study period was moved from August 1, 2022, to March 1, 2023. a 100-nurse non-probability (purposive) sample. Ten nurses who worked at Al-Basrah Teaching Hospital from September 6 to September 21 were the subjects of pilot research. Data were

\* Fundamentals of Nursing Department, College of Nursing, University of Basrah, Basrah, Iraq.

E-mail: ali.malik@uobasrah.edu.iq

\*\* College of Nursing, University of Basrah

gathered using the created questionnaire. Three sections make up the questionnaire: the first section covers sociodemographic information (age, gender, education level, working unit, and experience). A questionnaire with 26 questions about nurses' understanding of deep vein thrombosis makes up the second section. The survey's last portion includes seven inquiries about nurses' opinions on deep vein thrombosis. A panel of thirteen professionals evaluated the content of the tool for validity. On the basis of Cronbach's Alpha, the reliability of the questionnaire is assessed. The (SPSS ver26) was used to examine the data. Included in the study of descriptive data are the mean of score, standard deviation (SD), and frequency. A chi-square is a tool used in inferential data analysis. Each question in the second part consisted of two choices (True and False) and scored as (1 for a True answer and 0 for a False answer). By determining the cutoff threshold for the mean of the scores, the level of assessment for the second part's Knowledge scales was calculated and scored as follows: The researcher came up with the following numbers: (0-0.33) for poor Knowledge, (0.34-0.67) for moderate Knowledge, and (0.68-1) for good Knowledge. The Al-Basrah Health Director has obtained written official permissions to facilitate the data collection. Thirteen experts from College of Nursing/ University of Basrah selected to review the questionnaire.

## RESULTS

**Table 1:** The study sample's sociodemographic information

Characteristic	Classes	F	Percentage
Age	20 – 29	41	41%
	30– 39	18	18%
	40 – 49	24	24%
	50 and above	17	17%
Gender	Male	36	36%
	Female	64	64%
Education Level	Diploma	83	83%
	Bachelor's	17	17%
	Master's	0	0%
	Total	100	100%
Working unit	Surgical unit	53	53%
	Medical unit	47	47%
	Total	100	100%
Experience	<5 Years	36	36%
	5-10 Years	31	31%
	>10 Years	33	33%

**Table 2:** Nurses' Knowledge Regarding Deep Vein Thrombosis

Classification	Frequency	Percentage	Mean of Score	SD	Assessment
Poor	76	76 %	0.25	0.428	Poor
Moderate	22	22 %			
Good	2	2 %			
Total	100	100 %			

SD: standard deviation

**Table 3:** Nurses' Attitudes Regarding Deep Vein Thrombosis

Classification	Frequency	Percentage	Mean of Score	SD	Assessment
Poor	83	83 %	0.27	0.423	Poor
Moderate	13	13 %			
Good	4	4 %			
Total	50	100 %			

SD: standard deviation

## DISCUSSION

The study's findings indicate that the nurses' average age was between 20 and 29 years old, with a percentage (41%) of age being above 40. These outcomes supported According to the findings of a descriptive design study done at Al-Hilla Teaching Hospital by<sup>28</sup>, 83.0% of the sample was between the ages of 20 and 29. According to the current study's findings, women made up 64% of the nursing staff. The non-experimental descriptive design's findings that (90%) of nurses were female are supported by these findings<sup>29</sup>.The majority of nurses with diplomas (83%) graduated with a percentage, according to the study's findings. The results of the current study are consistent with those of a quasi-experimental study done by<sup>30</sup>, which discovered that 60% of nurses have diploma degrees. According to the study's results, 36% of nurses have less than five years of experience. These findings are consistent with the survey, which revealed that 43.37% of nurses had experience of between one and five years, whereas 30% had experience of more than a year and 53% had more. The results of the current investigation also showed that, compared to the (31) stated figure of 58%, 53% of nurses were working in the surgical unit. According to the investigation's findings, 76% of nurses don't know enough about deep vein thrombosis. Researchers think that many factors contribute to most nurses' lack of knowledge about deep vein thrombosis, including the fact that most of them have diplomas, that they didn't study the condition enough, that there aren't enough training programs for it, and that most of them don't actively pursue their education in the field. These results support the findings of a cross-sectional study by<sup>31</sup>, research discovered that most nurses know very little about the causes of DVT. These results refute those who<sup>1</sup> asserted that nurses were highly knowledgeable about deep vein thrombosis. According to the study's findings, 83% of nurses had negative attitudes concerning deep vein thrombosis. A study<sup>32-36</sup> that claimed that most nurses have positive attitudes towards deep vein thrombosis was at odds with the findings of the current investigation. The findings of this study showed a strong relationship between nurses' level of education and skill. These results corroborated a study<sup>37</sup> that discovered a strong relationship between nurses' expertise and level of education. The study's conclusions indicate that a nurse's level of knowledge is unrelated to their gender or number of years of experience. These conclusions were supported by a study<sup>38-40</sup> that discovered no connection between nurses' knowledge and their (gender and experience). The results of the current study also demonstrated that there are no significant differences in nurses' expertise based on their age or working unit. These results confirm the assertions made by those<sup>41</sup> who asserted that there are no discernible differences in nurses' knowledge based on their (age and working unit). The results of this study showed a strong relationship between nurses' attitudes and level of education<sup>42-47</sup>.

## CONCLUSION

1. The amount of education of nurses and their knowledge and attitudes are significantly correlated.
2. Age, gender, working unit, and years of experience of nurses do not significantly affect their knowledge and attitudes.

## RECOMMENDATIONS

1. Provide nurse education courses to help them learn more about deep vein thrombosis.
2. Offering nurses education programs to help them gain more insight into deep vein thrombosis.
3. Providing booklets for nurses related to deep vein thrombosis.
4. Due to a lack of studies on deep vein thrombosis in Iraq, the researcher suggested more research.

**Table 4:** Relationship between Socio-Demographic Characteristics of Nurses’ Knowledge toward Deep Vein Thrombosis

Socio-Demographic Characteristics	Classes	Knowledge			Significant
		Poor	Moderate	Good	
Age	20 – 29	28	12	1	Chi-square= 11.717 Df= 6 P-value= 0.069 NS
	30– 39	16	2	0	
	40 – 49	22	1	1	
	50 and above	10	7	0	
Gender	Male	30	6	0	Chi-square= 2.250 Df= 2 P-value= 0.325 NS
	Female	46	16	2	
Education Level	Diploma	73	8	2	Chi-square= 43.498 Df= 2 P-value= 0.000 HS
	Bachelor’s	3	14	0	
	Master’s	0	0	0	
Working unit	Surgical unit	41	12	0	Chi-square= 2.304 Df= 2 P-value= 0.316 NS
	Medical unit	35	10	2	
Experience	<5 Years	27	8	1	Chi-square= 2.238 Df= 4 P-value= 0.692 NS
	5-10 Years	26	5	0	
	>10 Years	23	9	1	

"df: Degree of freedom, NS: Not Significant, HS: High Significant"

**Table 5:** Association between Socio-Demographic characteristics of Nurses’ Attitudes toward Deep Vein Thrombosis

Socio-Demographic Characteristics	Classes	Attitudes			Significant
		Poor	Moderate	Good	
Age	20 – 29	30	9	2	Chi-square= 9.148 Df= 6 P-value= 0.165 NS
	30– 39	16	2	0	
	40 – 49	22	0	2	
	50 and above	15	2	0	
Gender	Male	30	4	2	Chi-square= 0.495 Df= 2 P-value= 0.781 NS
	Female	53	9	2	
Level of Education	Diploma	79	2	2	Chi-square= 53.937 Df= 2 P-value= 0.000 HS
	Bachelor’s degree	4	11	2	
	Master’s degree	0	0	0	
Working unit	Surgical unit	47	5	1	Chi-square= 2.800 Df= 2 P-value= 0.247 NS
	Medical unit	36	8	3	
Experience	<5 Y	29	5	2	Chi-square= 2.494 Df= 4 P-value= 0.646 NS
	5-10 Y	26	5	0	
	>10 Y	28	3	2	

**Authorship Contribution:** All authors share equal effort contribution towards (1) substantial contributions to conception and design, acquisition, analysis and interpretation of data; (2) drafting the article and revising it critically for important intellectual content; and (3) final approval of the manuscript version to be published. Yes.

**Potential Conflicts of Interest:** None

**Competing Interest:** None

**Acceptance Date:** 03-08-2023

**REFERENCES**

1. Al-Mugheed KA, Bayraktar N. Knowledge and practices of nurses on deep vein thrombosis risks and prophylaxis: A descriptive cross-sectional study. *J Vas Nurs* 2018;36(2):71-80.
2. Tiryag AM, Atiyah HH. Nurses' knowledge toward obesity in al-Basra city. *Annals of the Romanian Society for Cell Biology* 2021;4667-73.
3. Tiryag AM, Atiyah HH. Nurses’ Knowledge toward Bariatric Surgery at Surgical Wards at Teaching Hospitals in Al-Basra City. *Indian J Forensic Med Toxicol* 2021;15(3):5152-9.
4. Shahi A, Chen AF, Tan TL, et al. The incidence and economic burden of in-hospital venous thromboembolism in the United States. *J arthroplasty* 2017;32(4):1063-6.

5. Dai WL, Lin ZM, Shi ZJ, et al. Venous thromboembolic events after total knee arthroplasty: which patients are at a high risk?. *J Knee Surg* 2019;33(10):947-57.
6. Warren JA, Sundaram K, Kamath AF, et al. Venous thromboembolism rates did not decrease in lower extremity revision total joint arthroplasty from 2008 to 2016. *J arthroplasty* 2019;34(11):2774-9.
7. Warren JA, Sundaram K, Anis HK, et al. Have venous thromboembolism rates decreased in total hip and knee arthroplasty?. *J arthroplasty* 2020;35(1):259-64.
8. Gionis MN, Ioannou CV, Katsamouris AN, et al. The study of the thrombin generation mechanism and the effect of low molecular weight heparin as thromboprophylaxis in patients undergoing total knee and hip replacement. *Thromb Res* 2013;132(6):685-91.
9. Melinte RM, Arbănași EM, Blesneac A, et al. Inflammatory biomarkers as prognostic factors of acute deep vein thrombosis following the total knee arthroplasty. *Medicina* 2022;58(10):1502.
10. Reganon E, Vila V, Martínez-Sales V, et al. Sialic acid is an inflammation marker associated with a history of deep vein thrombosis. *Thromb Res* 2007;119(1):73-8.
11. Matos MF, Lourenço DM, Orikaza CM, et al. The role of IL-6, IL-8 and MCP-1 and their promoter polymorphisms IL-6-174GC, IL-8-251AT and MCP-1-2518AG in the risk of venous thromboembolism: a case-control study. *Thromb Res* 2011;128(3):216-20.
12. Hou H, Ge Z, Ying P, et al. Biomarkers of deep venous thrombosis. *J Thromb Thrombolysis* 2012;34(3):335-46.
13. Shi D, Xu X, Xu Z, et al. P-selectin: an unpredicted factor for deep vein thrombosis after total hip arthroplasty. *BioMed Res int* 2014;2014.
14. Xu Z, Shi D, Zhang C, et al. Postoperative plasma D-dimer value for predicting deep venous thrombosis following hip arthroplasty with nadroparin prophylaxis. *Hip Int* 2013;23(4):411-6.
15. Huang Z, Xie X, Li L, et al. Intravenous and topical tranexamic acid alone are superior to tourniquet use for primary total knee arthroplasty: a prospective, randomized controlled trial. *J Bone Joint Surg Am* 2017 Dec;99(24):2053-61.
16. Bottner F, Wegner A, Winkelmann W, et al. Interleukin-6, procalcitonin and TNF- $\alpha$ : markers of peri-prosthetic infection following total joint replacement. *J Bone Joint Surg Br* 2007;89(1):94-9.
17. Park KK, Kim TK, Chang CB, et al. Normative temporal values of CRP and ESR in unilateral and staged bilateral TKA. *Clin Orthop Relat Res* 2008;466(1):179-88.
18. Ettinger M, Savov P, Calliess T, et al. Improved diagnostic accuracy with the classification tree method for diagnosing low-grade periprosthetic joint infections by quantitative measurement of synovial fluid alpha-defensin and C-reactive protein. *Int Orthopaedics* 2020;44:31-8.
19. Bauer TW, Parvizi J, Kobayashi N, et al. Diagnosis of periprosthetic infection. *J Bone Joint Surg* 2006;88(4):869-82.
20. Cheng X, Zhang M, Xie Y, et al. Bone marrow-derived mesenchymal stem cells accelerate angiogenesis in pregnant experimentally induced deep venous thrombosis rat model via up-regulation of pro-angiogenic secretogranin II. *Int Immunopharmacol* 2023;118:110025.
21. Yohannes S, Abebe T, Endalkachew K, et al. Nurses' knowledge, perceived practice, and their associated factors regarding Deep Venous Thrombosis (DVT) prevention in Amhara Region Comprehensive Specialized Hospitals, Northwest Ethiopia, 2021: a cross-sectional study. *Critical Care Res Prac* 2022;2022.
22. Barco S, Woerschling AL, Spyropoulos AC, et al. European Union-28: an annualised cost-of-illness model for venous thromboembolism. *Thromb Haemost* 2016;115(04):800-8.
23. Mulatu A, Melaku T, Chelkeba L. Deep venous thrombosis recurrence and its predictors at selected tertiary hospitals in Ethiopia: a prospective cohort study. *Clin Appl Thromb Hemost* 2020;26:1076029620941077.
24. Yu-Fen MA, Yuan XU, Chen YP, et al. Nurses' objective knowledge regarding venous thromboembolism prophylaxis: A national survey study. *Med* 2018;97(14).
25. Lee JA, Grochow D, Drake D, et al. Evaluation of hospital nurses' perceived knowledge and practices of venous thromboembolism assessment and prevention. *J Vas Nurs* 2014;32(1):18-24.
26. Shaaban AE. Effect of nursing care protocol on deep vein thrombosis occurrence among critically neurological patients. *Port Said Scientific J Nurs* 2021;8(1):206-25.
27. Galanaud JP, Sevestre MA, Genty C, et al. Incidence and predictors of venous thromboembolism recurrence after a first isolated distal deep vein thrombosis. *J Thromb Haemost* 2014;12(4):436-43.
28. Aydin AK, Karadag A. Assessment of nurses' knowledge and practice in prevention and management of deep tissue injury and stage I pressure ulcer. *J Wound Ostomy Continence Nurs* 2010;37(5):487-94.
29. Mohammed AS, Taha NM, Abdel-Aziz EM. Nurses' performance regarding venous thromboembolism prophylaxis at intensive care unit. *Zagazig Nurs J* 2018;14(1):1-7.
30. Elkattan BE, Elderiny SN. Effect of nursing care guidelines on preventing deep venous thrombosis among patients undergoing arthroplasty surgery. *International J Novel Res Healthcare Nurs* 2017;6(2):757-74.
31. Parveen B, Kousar Parveen MH, Afzal M. Nurses Knowledge And Practices Regarding Risk Factors And Prevention of Patients Diagnosed with Deep Vein Thrombosis. *J Global Biosci* 2021;10(6):8780-800.
32. Yan T, He W, Hang C, et al. Nurses' knowledge, attitudes, and behaviors toward venous thromboembolism prophylaxis: how to do better. *Vascular* 2021;29(1):78-84.
33. Collins R, MacLellan L, Gibbs H, et al. Venous thromboembolism prophylaxis: The role of the nurse in changing practice and saving lives. *Australian J Adv Nurs* 2010;27(3):83-9..
34. Sahu R, Menon SA, Anik S, et al. A study to assess the effectiveness of planned teaching programme on knowledge regarding Deep Vein Thrombosis among B. Sc. Nursing intern's working in Jawaharlal Nehru Hospital and Research Centre Bhilai (CG). *Int J Nurs Educ Res* 2017;5(3):272-8.
35. Younis NM, Taher AK. Efficacy of Trans Theoretical Model Intervention for Improving Behaviors related to Electronic Hookah Smoking among Healthcare Workers in Mosul Hospital: A Randomized Control Trail. *Int J Membrane Sci Tech* 2023;10(2):1433-9.
36. Ali HA, Abbas FF, Younis NM. Mothers' knowledge and attitudes towards breastfeeding in Thi-Qar City, Iraq. *Rawal Med J* 2023;48(2):514.
37. Bura'a LN, Younis NM. Nurses knowledge regarding to phototherapy at neonatal care units in Mosul City, Iraq. *Rawal Med J* 2023;48(2):379.
38. Ahmed M M, Naji A B, Younis N M. Efficacy of an educational program based on health belief model to enhancing weight control behaviors among employees in the University of Mosul: a randomized controlled trial. *Revis Bionatura* 2023;8 (3) 28.
39. Younis NM. Evaluation the health lifestyle of kindergarten students at Mosul city/Iraq. *Int J Med Toxicol Legal Med* 2023;26(1):148-52.
40. Bura'a LN, Younis NM. An Interventional Program on Nurses Knowledge and Practice towards Phototherapy in Neonatal Care Units. *Int J Membrane Sci Tech* 2023;10(2):1428-32.
41. Al-Snafi AE, Alfuraiji N. Medicinal Plants with Anti-Obesity Effects: A Special Emphasis on Their Mode of Action. *Bahrain Med Bull* 2023;45(2).
42. Ayed AY, Younis NM, Ahmed MM. Comparison of infection severity of vaccinated and unvaccinated health workers with Corona Virus: A cohort study. *J Edu Health Promotion* 2023;1(1):336.

43. Mahdi NS, Mansour KA. Nurses' Knowledge toward Preventive Measure of Pulmonary Embolism in AL Nasiriyah City. *Annals of the Romanian Society for Cell Biology* 2021 29;25(7):497-506.
44. Younis NM. Epidemiology of Hepatitis B-virus in Nineveh province: Retrospective Study. *Int J Membrane Sci Tech* 2023;10(2):1440-4.
45. Asiri NS, Alshehri AA, Alshehri FF. Knowledge, Attitude and Practice of First Aid Management of Epistaxis Among General Population in Saudi Arabia. *Bahrain Med Bull* 2023;45(1).
46. Younis NM. Prevalence of Electronic Hookah and Risk Factors among University Students in Mosul City/Iraq. *Int J Membrane Sci Tech* 2023;10(2):1422-7.
47. Tiryag AM, Atiyah MA, Khudhair AS. Nurses' Knowledge and Attitudes toward Thyroidectomy: A Cross-Sectional Study. *Health Edu Health Promotion*. 2022;10(3):459-65.