

## **The Evaluation of Not Stenting after Uncomplicated Ureteroscopy: A Randomized Prospective Study**

**Waleed Ali, Arab Board Part (A)\* Reem Al-Bareeq, MD\*\*  
Moh'd R.Samiei, FRCS\*\*\* Shaker Al-Muttawa, Facharzt, Arab Board(Surg)\*\*\***

**Objective:** The aim of this study is to determine the morbidity associated with not placing a ureteral stent following uncomplicated ureteroscopy and to determine the safety of performing ureteroscopy as a day-case procedure.

**Design:** This is a prospective study.

**Setting:** The study was done in the Day-case Unit, Salmaniya Medical Center, Bahrain.

**Methods and Materials:** Sixty-one patients with symptomatic ureteric stones who underwent ureteroscopy as an outpatient procedure. The patients were followed up at days 1-4 and again at 4-6 weeks postoperatively to determine the morbidity associated with the procedure (i.e. analgesic use and length of time to return to normal activity). Follow up consisted of urine analysis and radiological studies, which were done between 2-3 months postoperatively.

**Results:** Twenty one patients (34%) had no discomfort postoperatively. The remaining 40 patients (66%) had discomfort, 35 (57%) had pain that was relieved by the 2nd postoperative day. The main indication for admission was pain and hematuria. Five patients required admission. None of the patients had major immediate postoperative complications and all of them had follow up excretory urography or ultrasound done. None showed newly identified ureteric strictures or stone recurrence.

**Conclusions:** Day-case ureteroscopic management of ureteric stones is a potentially safe procedure and can be performed in selected patients. We believe that routine placement of ureteric stents following uncomplicated ureteroscopy is not essential.

*Bahrain Med Bull 2001;23(1):34-36.*

---

\* Senior Urology Resident

\*\* Junior Resident

\*\*\* Senior Urology Consultant

Department of Surgery

Salmaniya Medical Complex

State of Bahrain

Ureteroscopy is one of the major endourologic procedures used for diagnostic and therapeutic purposes. The main indication for this procedure is ureteric stones, but other indications would include tumor lesions of the ureter, renal pelvis and ureteral strictures<sup>1</sup>. This procedure is usually performed as an in-patient procedure in Bahrain since most patients are admitted through the Emergency department. However, in North America and other centers, it is mainly performed on an outpatient basis<sup>1</sup>. More centers worldwide are taking on that protocol, which is more efficient and cost-effective. The routine placement of a ureteral catheter or stent following ureteroscopic stone manipulation is widely recommended<sup>2</sup>. The rationale for stent placement is to reduce or prevent postoperative ureteric strictures and to prevent colic secondary to postoperative ureteral edema<sup>3</sup>.

This randomized, prospective study aims to determine the safety of performing ureteroscopy as a day case procedure and to determine the morbidity associated with not placing a stent following uncomplicated ureteroscopy.

## **METHODS**

From July 1997 to December 1999, we have prospectively followed 61 consecutive patients who underwent ureteroscopy without stenting for the management of ureteric stones. Forty were male with a mean age of 26 years and 21 were female with a mean age of 21 years. One patient was excluded from the study because we lost him for follow up.

The patients were treated on an outpatient basis. The procedure was performed under general anesthesia. Balloon ureteric orifice dilatation was performed in all patients. A Wolf 8.5Fr rigid ureteroscope was introduced and the stone was fragmented with a Swiss lithoclast pneumatic lithotripter.

Postoperatively all patients received one dose of an intramuscular narcotic analgesic (pethidine) and were provided with a Dextropropoxyphene + Paracetamol (Distalgesic) prescription.

The patients were informed that there would be some discomfort at the end of the procedure.

The patients were followed up between 1-4 days and a second time between 4-6 weeks postoperatively. They were asked whether they had visited the emergency room for any post-operative complications, their use of analgesics and the time required to return to normal activity. Patients were asked to return for a radiological follow up with a renal ultrasound and excretory urogram postoperatively.

## **RESULTS**

Fifty-five patients (90%) had pain at presentation, 61 (100%) had microscopic hematuria, 40 (66%) had dysuria and 40 (66%) had increased frequency of micturation.

All the patients had a KUB preoperatively, while ultrasound was done for 13 (21%) patients and IVP was performed for 48 (79%).

The main indication for surgery was pain and/or obstruction.

Eleven (18%) patients had upper ureteric stones, while 50 (82%) had lower ureteric stones. The stone size ranged from 4 mm to 1.8 cm. About 14 (23%) patients had multiple stones and 47 (77%) patients had single stones.

Twenty-one patients (34%) had no post-operative pain. Forty patients (66%) had pain which lasted up to the time they were kept in the recovery room (Table 1).

**Table 1. Duration of Post-operative Pain**

<b>Duration of Post-operative Pain</b>	<b>No. of Patients</b>
No pain	21 (34%)
Recovery Room (4-6hrs)	40 (66%)
Within 24hrs	26 (43%)
1 <sup>st</sup> Post-op Day	16 (26%)
2 <sup>nd</sup> Post-op Day	5 (8%)
Later	Nil

Only 5 (8%) patients required hospitalization, 4 (7%) were females and one was a male patient. The reason for their admission was severe pain and hematuria immediately after the operation (Table 2).

**Table 2. Number of patients with Post-operative complications**

<b>Complications</b>	<b>No. of Patients</b>
Pain and hematuria	5
Recurrence	Nil
Ureteral stricture	Nil

There were no major complications in the immediate postoperative period.

All patients were followed up with ultrasound and IVP. No disorders were detected except one patient was re-operated upon for a stone on the contralateral side. We lost track of one patient in the follow-up period.

## **DISCUSSION**

Endoscopic procedures mainly ureteroscopy, are performed as day case surgeries in most of North American urologic centers. Some authors have even reported performing such surgeries with intravenous sedation in a large number of out-patient series<sup>4</sup>. In Bahrain, this type of surgery is mainly done as an inpatient procedure.

Although ureteric stenting after ureteroscopy has been recommended to prevent the development of ureteric strictures and relieve any ureteral edema<sup>1,5</sup>, there have been no

controlled trials in humans to our knowledge which have shown that it prevents strictures.

There are several reports suggesting that stent placement may be associated with significant symptoms such as chronic infection, irritation and stone formation and these symptoms persist until the stent is removed<sup>6-8</sup>. We are not aware of any comparative studies of patients with and without stents following ureteroscopy, or any reports suggesting any symptomatic advantage gained by placing a stent.

All 61 patients in this prospective study had a KUB done to establish the diagnosis of ureteric calculus which were mainly situated in the lower ureter. The majority of them presented with microscopic hematuria. These patients were selected to undergo ureteroscopy as an outpatient without stent placement.

Follow-up of the patients and assessment of the duration of post-operative pain revealed that 34% were pain free. The majority of patients (66%) had pain up to 4-6 hrs in the recovery room. None had continuous pain beyond the 2<sup>nd</sup> post-operative day (Table 1). There were five patients who required admission due to pain (3patients), pain and hematuria (2 patients). They were discharged after the first post-operative day (Table 2).

Most patients required one dose of intramuscular narcotic (Pethidine) post-operatively. Patients used oral analgesics to relieve the pain.

Further follow-up at 4-6 weeks revealed no incidence of ureteric stricture or recurrence of the calculus in any of the 61 cases.

Although traditional protocol necessitates the placement of a stent, various studies have showed its irrelevance in uncomplicated ureteroscopies. Hosking et al reported a good outcome in 93 patients who underwent ureteroscopy without stent placement as an outpatient procedure. About 43% had no discomfort and the remaining had discomfort that was short-lived ( $\leq 2$ days)<sup>9</sup>. Similarly, our study showed that the majority of patients had discomfort which did not last more than 4-6 hours in the recovery room, it was controlled with one dose of intravenous narcotic post-operatively. This supports the idea that ureteroscopy done for ureteral calculi can successfully be managed in the outpatient unit and does not need the placement of a stent if the procedure is uncomplicated.

A potential problem associated with this approach is predicting those patients who might develop severe pain and hematuria, which demands admission regardless of an uneventful procedure, although they constitute a small number of patients. Rane et al experienced only one admission essentially for loin pain after ureteroscopy as an outpatient procedure without stenting<sup>10</sup>. Although the procedures were uneventful, five patients in our study required admission. There were no obvious indicators to predict that these patients would potentially need subsequent admission. To manage this problem, experienced nursing staff, immediate follow-up and an adequate recovery time should be allowed, so as to determine which patients will need admission post-operatively.

We do believe that routine placement of stents following complicated ureteroscopy is

indicated, but for straightforward uncomplicated procedures it is not recommended.

## CONCLUSIONS

**The results of this study indicate that ureteroscopy performed in the outpatient unit can be a safe alternative to traditional hospital admission in selected patients. Ureteric stenting is not a necessary procedure after ureteroscopy as long as it is uneventful. However, close post-operative observation is essential to determine those small number of patients who require admission post-operatively due to minor complications (pain and hematuria).**

## REFERENCES

1. Harmon WJ, Sershon PD, Blute ML, et al. Ureteroscopy: current practice and long-term complications. *J Urol* 1997;157:28.
2. Bolton DM. Techniques in rigid ureteroscopy. In: Smith's Textbook of Endourology. Smith AD, Badlani GH, Bagley DH, et al, eds. St. Louis: Quality Medical Publishing Inc, Vol 1, Chapt 31 1996: 435-42.
3. Weinberg, JJ, Synder JA, Smith AD. Mechanical extraction of stones with rigid ureteroscopes. *Urol Clin N Am* 1998;15:339.
4. Hosking DH, Bard RJ. Ureteroscopy with intravenous sedation for treatment of distal ureteric calculi: a safe and effective alternative to shock wave lithotripsy. *J Urol* 1996;156: 24.
5. Boddy SM, Nimmon CC, Jones S, et al. Acute ureteric dilatation for ureteroscopy-an experimental study. *Br J Urol* 1988;61:27.
6. Pryor JL, Langley MJ, Jenkins AD. Comparison of symptom characteristics of indwelling ureteral catheters. *J Urol* 1991;145:719.
7. Pollard SG, Macfarlane R. Symptoms arising from Double-J ureteral stents. *J Urol* 1988;139:37.
8. Bregg K, Riehle RA. Morbidity associated with indwelling internal ureteral stents after shock wave lithotripsy. *J Urol* 1989;141:150.
9. Hosking DH, McColm SE, Smith WE. Is stenting following ureteroscopy for removal of distal ureteral calculi necessary? *J Urol* 1999;161:48-50.
10. Rane A, Cahill D, Larner T, et al. To stent or not to stent? That is still the question. *J Endourol* 2000;14:479-81.